

CHORIOAMNIONITIS - SS

NURSING

- In and Out straight catheterization: Utilize collection for any urine studies ordered.

IV FLUIDS

- Lactated Ringers IV to run at 125 mL/Hr
- Sodium Chloride 0.9% IV to run at 125 mL/Hr

MEDICATIONS

Antipyretic medications

- Acetaminophen (Tylenol) 650 mg Po Q 4 Hrs Prn Temp > 101 °F.
Total Acetaminophen not to exceed 4,000 mg/24 Hrs

Antibiotic medications

- Ampicillin (Omnipen) 2 Gm IVPB Q 6 Hrs
- Clindamycin (Cleocin) 900 mg IVPB Q 8 Hrs
- Gentamicin IV per Pharmacy protocol. Indication: Chorioamnionitis
- MetroNIDAZOLE (Flagyl) 500 mg IVPB Q 8 Hrs
- Vancomycin (Vancocin) IV per Pharmacy protocol to achieve target trough levels of 10 – 15 mg/L.
Indication: Chorioamnionitis

Other medications: _____

***All labs/diagnostics will be drawn/done routine now unless otherwise specified**

LABORATORY- Hematology

- Complete Blood Count (CBC)

LABORATORY- Chemistry

- Basic Metabolic Panel (BMP)

LABORATORY- Urine

- Urinalysis Reflex Culture: Type of Specimen: Catheterized

MICROBIOLOGY

- Culture, Blood (BC) X 1

MD CONSULTS

Reminder: Consider specialty referral (Neonatology, Maternal Fetal Medicine)

- Consult MD _____
- Consult MD _____