

TRANSFUSION NOW SS - INPATIENT

BLOOD BANK / NURSING

****SELECT the order for the blood product. In addition fill out the corresponding TRANSFUSION INSTRUCTION order (below) for nursing to administer the blood product*****

Red Blood Cells

PRBC _____ **unit(s)**

PRBC Indications

- | | | |
|--|--|--|
| <input type="checkbox"/> Hemoglobin <8 g/dL | <input type="checkbox"/> Preop Hgb <9 g/dL | <input type="checkbox"/> Blood Loss >15% |
| <input type="checkbox"/> Symptomatic Anemia | <input type="checkbox"/> Hypotension/Shock | <input type="checkbox"/> Chronic Renal Failure |
| <input type="checkbox"/> Chemotherapy <input type="checkbox"/> Other _____ | | |

Unit specification

- | | | | | |
|-------------------------------------|-----------------------------------|-------------------------------------|---------------------------------------|-------------------------------------|
| <input type="checkbox"/> Autologous | <input type="checkbox"/> Directed | <input type="checkbox"/> Irradiated | <input type="checkbox"/> CMV Negative | <input type="checkbox"/> Washed RBC |
|-------------------------------------|-----------------------------------|-------------------------------------|---------------------------------------|-------------------------------------|

PRBC Transfusion Instructions:

Begin Transfusion At: _____

Transfuse during hemodialysis yes no

Additional Instructions: _____

Platelets

Platelets Apheresis _____ **unit(s)**

Platelet Indications

- | | | |
|--|--|---|
| <input type="checkbox"/> Plat<20,000 u/L | <input type="checkbox"/> Plat<50,000 u/L surgery pending | <input type="checkbox"/> Platelet Dysfunction |
| <input type="checkbox"/> Preop | <input type="checkbox"/> Hypotension/Shock | <input type="checkbox"/> Blood Loss |
| <input type="checkbox"/> Chemotherapy <input type="checkbox"/> Other _____ | | |

Unit specification

- | | |
|-----------------------------------|---------------------------------------|
| <input type="checkbox"/> Directed | <input type="checkbox"/> CMV Negative |
|-----------------------------------|---------------------------------------|

Platelets Transfusion Instructions:

Begin Transfusion At: _____

Additional Instructions: _____

Thawed Plasma

Thawed Plasma _____ **unit(s)**

Thawed Plasma Indications

- | | | |
|---------------------------------------|--|--|
| <input type="checkbox"/> INR >1.5 | <input type="checkbox"/> Coumadin reversal | <input type="checkbox"/> Hypotension/Shock |
| <input type="checkbox"/> Preoperative | <input type="checkbox"/> Blood Loss | <input type="checkbox"/> Chemotherapy |
| <input type="checkbox"/> Other _____ | | |

Plasma Transfusion Instructions:

Begin Transfusion At: _____

Additional Instructions: _____

Cryoprecipitate

Cryoprecipitate _____ **unit(s)**

Cryoprecipitate Indication

- | | | |
|--|---|--|
| <input type="checkbox"/> Fibrinogen <100 mg/dL | <input type="checkbox"/> Von Willebrand Disease | <input type="checkbox"/> Hypotension/Shock |
| <input type="checkbox"/> Blood Loss | <input type="checkbox"/> Chemotherapy | <input type="checkbox"/> Other _____ |

Cryoprecipitate Transfusion Instructions:

Begin Transfusion At: _____

Additional Instructions: _____

Granulocytes

Granulocytes _____ **unit(s)**

Granulocytes Transfusion Instructions

Additional Instructions: _____

MEDICATIONS

- Sodium Chloride 0.9% 100 mL IV x 1 dose for priming/post reaction flushing.

Acetaminophen premed orders

- Acetaminophen (Tylenol) 650 mg Po x 1 dose as a pre-medication 30 mins prior to starting PRBC, FFP, or Platelet transfusions. *Total Acetaminophen not to exceed 4,000 mg/24 Hrs*
- Acetaminophen (Tylenol) 650 mg PR x 1 dose as a pre-medication 30 mins prior to starting PRBC, FFP, or Platelet transfusions. Give if pt unable to take Po Acetaminophen (if ordered). *Total Acetaminophen not to exceed 4,000 mg/24 Hrs*

DiphenhydrAMINE premed orders

- DiphenhydrAMINE (Benadryl) 25 mg Po x 1 dose as a pre-medication 30 mins prior to starting PRBC, FFP, or Platelet transfusions.
- **For Patients > or = 65 years old unable to take Po meds, use 12.5 mg order**
- DiphenhydrAMINE (Benadryl) 12.5 mg IV Push x 1 dose as a pre-medication 30 mins prior to starting PRBC, FFP, or Platelet transfusions. Give if pt unable to take Po DiphenhydrAMINE (if ordered).
- **For Patients < 65 years old unable to take Po meds, use 25 mg order**
- DiphenhydrAMINE (Benadryl) 25 mg IV Push x 1 dose as a pre-medication 30 mins prior to starting PRBC, FFP, or Platelet transfusions. Give if pt unable to take Po DiphenhydrAMINE (if ordered).

Furosemide orders

- Furosemide (Lasix) 20 mg IV Push x 1 dose after 1st unit of PRBC.
- Furosemide (Lasix) 40 mg IV Push x 1 dose after 1st unit of PRBC.
- Furosemide (Lasix) 20 mg IV Push x 1 dose after 3rd unit of PRBC.
- Furosemide (Lasix) 40 mg IV Push x 1 dose after 3rd unit of PRBC.